

## Toxicological Effects of Aqueous Extract of *Calotropis procera* Leaves in Experimentally Poisoned Rabbits

Ahmad H. Al-Zuhairi<sup>1\*</sup>, Jenan M. Khalaf<sup>2</sup>, and Salah N. Ibrahim<sup>3</sup>

<sup>1</sup>Department of Internal and Preventive Medicine, College of Veterinary Medicine, University of Diyala,

<sup>2</sup>Department of Internal and Preventive Medicine, College of Veterinary Medicine, University of Baghdad, Iraq,

<sup>3</sup>Department of Physiology and Pharmacology, College of Veterinary Medicine, University of Diyala, Iraq

### ABSTRACT

*Calotropis procera* is announced to be medicinal and poisonous plant to human and animals. In this study, the toxicological effects of aqueous leaves extract were evaluated experimentally in rabbits. The median lethal dose (LD<sub>50</sub>) was estimated at 2435.25 mg/kg BW using the Up and Down method. Twenty-five local breed rabbits, 1-2-year-old and 1-1.9 kg BW, were divided randomly into five groups with five rabbits each. Those of groups I, II, III and IV were exposed to 1/10 of LD<sub>50</sub> (243.5mg/kg), 1/12.5 of LD<sub>50</sub> (194.8 mg/kg), 1/15 of LD<sub>50</sub> (162.4 mg/kg), and 1/20 of LD<sub>50</sub> (121.8 mg/kg BW of extract respectively for 8 weeks, while those in G.V were left without exposed as a control group. Alkaloids, saponins, tannins, cardiac glycosides, steroidal, glycoside, terpenoids and flavonoids detected in phytochemicals screening. The dependent parameter were clinical signs exhibited by animals during the study in addition to some hematological parameters Red blood cells (RBC) count, hemoglobin (Hb) concentration and packed cell volume (PCV). The main changes observed during monitoring the animals were chewing of mouth, bradycardia, engorged blood vessels, coughing and depression especially in these of G I, II and III; diminution body weight in G I; abortion in G I and II. The hematological and biochemical parameters depended showed a significant increase during the study. From this we can concluded that aqueous extract of leaves of the plant has a harmful effect in rabbits.

**Keywords:** *Calotropis procera*, Rabbits, Toxicity, Hematological, Biochemical parameters

### Introduction

Poisonous plants are among the important causes of economic losses to livestock (1). Family of Asclepiadaceae include genus *Calotropis* which have two species *C. procera* and *C. gigantea*; while *C. procera* it is a soft wooded, perennial, evergreen bush. distributed in all parts of the world at a hot weather, dry and alkaline ground (2-4). *Calotropis procera* famous by different vernacular nouns as Giant milkweed and Swallow wort in English, debaj, usher, oshar in Arabic; Indian milkweed and

madar in Hindi ; bomba, algodónextranjero, cazuela in Spanish, french cotton in French; *Calotropis*, rubber bush, rubber plant, King Edward's crown, Prince of Wales' crown, kapok tree, king's crown kapok and others (4-8). Leaves of oshar screened to determine phytochemical compounds by many investigators, results revealed presence of mixtures of cardenolides, glycosides, phenols, tannins, saponins and terpenoids (9-11). Giant milkweed is well known for its medical as well as venomous characteristic (12). All parts of plant used medicinally to medication common diseases as anti-inflammatory, anti-pyretic, anti-parasites and anti-microbial activities, analgesic, cold, indigestion, diarrhea, eczema and rheumatism. It has been applied for treated many other conditions including, epilepsy, cancer, and snake bites, the smoke from the bark has been inhaled to treated of asthma and coughs (8).

Regardless these employ, *C. procera* can cause varying venomous influence in human being and

\*Correspondence: [ahmhansh@gmail.com](mailto:ahmhansh@gmail.com), Department of Internal and Preventive Medicine, College of Veterinary Medicine, University of Diyala, Iraq.

Received: 27 November 2019, Accepted: 22 April 2020, Published: 28 June 2020.

This article is an open access article under the terms and conditions of the Creative Commons Attribution License (CC BY 04 <https://creativecommons.org/licenses/by/4.0>).

DOI: <https://doi.org/10.30539/ijvm.v44i1.934>

animals via touch, air borne allergies, and consuming livestock to this tree (13). The whole plant (leaves, bark, stem latex and roots) is toxic. Brazilian farmers suggested has been accidental ingestion of fresh leaves of Giant milkweed as a toxic to ruminants by several semiarid regions.

Sheep consumption to swallow wort leaves is responsible of temporary cardiac arrhythmia at auscultation 30 min after dosage. *C. procera* plant is poisonous to heart and liver (11). In addition to hepatic poisoning, other investigators to notarize the renal toxicity of this plant (14,15). To best illustrate of the venomous influence of *C. procera*, a toxicological evaluation of the aqueous extract of leaves of this plant was take on clinical, hematological and biochemical parameter.

## Materials and Methods

### Experimental Design

This study was carried out after approval of the scientific committee of department of Internal and Preventive Medicine, College of Veterinary Medicine, University of Baghdad and accordance the International Standards of animal welfare.

### Collection and Preparation of Plant Material

Fresh leaves of *C. procera* were shear in July 2018, from villages of Almuqdadiya city, Diyala province, Iraq. Reference sample was identified by Ministry of Agriculture/ Directorate of Seeds Testing and Certification (D.S.T.C) in Abu Graib /Baghdad, (NO. document:41, at 7/1/2019). The fresh leaves were washed in tap water, dried in shade at room temperature, the dried leaves were grinded by electrical mill and this powder preserved at sealed box till used after that.

### Extraction of the Plant Material

The dried ground plant (50 g) was mixed with 2 L of deionized water, left for 72 hours in room temperature according to (16). The aqueous extract was filtrated using by Whatman filter paper, then evaporated on water bath to a dark green gummy residue.

### Acute Toxicity of *C. procera*

The estimation of oral LD<sub>50</sub> in male albino rabbits was done using the Up and Down method according to (17).

### Experimental Animals

Twenty five adult male and female rabbits were randomly distributed into five sets (I, II, III, and IV) daily exposed to (243.5, 194.8, 162.4, 121.8) milligram/kilogram of b.w respectively of leaves extractor of *Calotropis procera* for eight weeks and group V serving as the control. The dependent parameter was clinical examinations, including pulse and respiratory rate, body temperature, and body weight, in addition to monitoring for any abnormal changes in behavior, appetite, or other signs.

### Blood Sample Collection

Four milliliter of blood samples were obtained via cardiac puncture at a zero, 4 and 8 weeks, to study hematological and biochemical parameters, first part of blood sample was directly put into a tube containing (EDTA) as an anticoagulant. For biochemical analysis the second part of blood was placed in a tube without anticoagulant and centrifuged at 3000 round per minute for 5 minutes. The sera obtained were preserved at -4°C prior to analysis (18).

### Hematological Study

Red and white blood cells (RBC, and WBC) count was assess by Coles method (18). Hemoglobin (Hb), as well as, Packed cell volume (PCV) were determined by the Hb Hemoglobin Test Strips method.

### Biochemical Analysis

Glutamic-oxaloacetic transaminase (GOT) and glutamate pyruvate transaminase (GPT) were estimated as described by commercial kit SGOT, SGPT from AGAPPE DIAGNOSTICS SWITZERLAND, Alkaline phosphatase (ALP) by BIOBO, France. Level of serum creatinine was evaluated by creatinine kinetic technique kit BIOLABO, France, while serum urea level was estimated by using commercially available kit (UREA Berthelot from LINEAR CHEMICALS S.L).

### Preliminary Qualitative Phytochemical Analysis

*Calotropis procera* extract was submitted to phytochemical checking according to standard

procedure for each test, to determine for the existence of the secondary metabolites; alkaloids according to (19). Glycosides, tannins, and saponin components was carried out according to Harborne (20), steroids and terpenoids components, this detection was conducted according to (21). Cardiac glycosides detected according to (22). Detection of flavonoids components, this process was prepared according to (23).

### Statistical Analysis

The datum collected were analyzed employed by analyses of variance (ANOVA, 2 way), according to (24). Quantitative data are offered at a mean with standard error mean (SEM). Statistically significant was deemed to be at  $P \leq 0.05$ .

### Results and Discussion

$LD_{50} = xf + kd = 2250 + (0.741) \times 250 = 2435.25$  mg/kg BW. The plant used in current study is

*Calotropis procera* according to Ministry of Agriculture/ Directorate of Seeds Testing and Certification (D.S.T.C) in Abu Graib /Baghdad. Phytochemical checking of the leaves of *Calotropis procera*, by more than way in some tests determined the existence of flavonoids, alkaloids, terpenoids, steroids, saponins, tannins, glycosides and cardiac glycosides (Table1), which are known to possess medicinal and pesticidal properties (25). This result is in agreement with previous study (26).

### Acute Toxicity Study of Plant Extracts

Determination of median lethal dose ( $LD_{50}$ ) were conducted in male rabbit by up and down method, according to (17), the value of  $LD_{50}$  was 2435.25 mg/kg BW (Table 2).

**Table 1. Phytochemical analysis of leaves of *Calotropis procera***

Test	<i>C. pleaves powder</i>	Results
Flavonoid	++	Yellow Color
Alkaloid / Mayer's T.	++	White Precipitate
Terpenoid	++	Brown Color
Steroid	+++	Blue-Green Color
Saponins/ Stirring T.	+++	Big Foam
Saponins/ Silver Nitrate T.	++	Silver Mirror
Tannins/ Lead Acetate T	++	Gelatin-White Color
Tannins / Ferric chloride T.	++	Blue –Green Color
Glycosides/ Fehling,T.	++	Red Precipate
Glycosides/ Benedict s T.	–	Red Precipate
Cardiac glycosides	++	brown ring

T=test, += low concentration, ++= medium concentration, +++=high concentration, - =not detectable

**Table 2.  $LD_{50}$  of aqueous extract of Leaves of *Calotropis procera***

Range of doses	Decrease or increase in doses	Death or survival of animals within 24 hours	Value of K Table	Last used dose (xf)	Value of $LD_{50}$
1250-2500 mg/kg BW	250 mg/kg BW	OOOOXOXO	0.741	2250 mg/kg BW	2435.25 mg/kg BW

O=Survival, X=death

## Sub chronic Toxicity Study

### Clinical Signs

The main clinical signs appeared on animals through daily observation were as shown in Table 3. No signs appeared for first week. While in second week, those of group I (G I) were showed signs of convulsion, chewing movement of mouth, aching of ear and extremities, followed by depression, abnormal behavior included bite of legs, back aching chewing foreign materials around them. At 9<sup>th</sup> day vaginal discharge with abortion of 4 fetuses in 2<sup>nd</sup> day, and 17<sup>th</sup> day other rabbit aborted. In 6<sup>th</sup> week two rabbits died, one of them was suffered from diarrhea, severe depression, anorexia, stand in water, unable to stand for two days, then died.

The signs appeared on those from groups I and II continued with appearance of sneezing, coughing, diarrhea with depression. At 6<sup>th</sup> week animals in G I and group II (G II) showed engorgement of blood vessels of ears. Two animals in G II were aborted in 20<sup>th</sup> and 24<sup>th</sup> day, in 7<sup>th</sup> week one rabbit was suffered from depression and anorexia for three days, then died. The abnormal heartbeat and bradycardia are a signal to some harm hit on the heart, and this can be referring to cardiac glycosides effect. Appearance the blood vessels of ears engorged, that refers to circulatory disorder. disturbance of heart architecture, with popularize necrosis of the myocardium may be refers to cardiac glycosides which have the capability to curb of the membrane potential via crippling the Na /K ATPase pump (27).

Inhibition of the Na-K-ATPase in cardiac and other tissues by Cardiac glycosides, leading to intracellular detention of Na, pursued by rise intracellular Ca<sup>2+</sup> concentrations through the influence of the Na -Ca<sub>2</sub> exchange. The increased intracellular Ca<sub>2</sub> concentration boost inotropic and bradycardia, (28-30). Diarrhea and loss of appetite may be as a result of erosion of mucous lining of the gastrointestinal tract, similar result recorded by (31). Existence of the cardiac glycosides, tannins and alkaloids in several plant promote adverse influence regard to livestock (32), and Saponins disrupt cellular membranes (33). Faye, (1985) (34) revealed that the *C. procera* is a highly toxic when consummated with feed, resulting in high abortion rates .Similar result of abortion in current study, revealed by (35) , while (36) observed that *C. procera* has uterine stimulating effect. Body weight

of animals in all groups were non significantly increased except in those of GI. These results are in agreement with data observed by (26).

Heart rates of those groups (I, II and III) were decreased significantly with no significant difference in groups IV and V. Respiratory rates increased significantly in those of GI only. No significant changes in body temperature in all groups (Table 4). The results of RBC in current study did not should any significant changes in values of total erythrocyte counts in all groups during 4<sup>th</sup> weeks, and significantly increased in G I only during 8<sup>th</sup> weeks (Table 5). Value of hemoglobin concentration was changed in GI during 4<sup>th</sup> weeks and changed in GII and G IV during 8<sup>th</sup> weeks (Table 5). Values of PCV were significantly increased in GI during 4<sup>th</sup> and 8<sup>th</sup> weeks and during 8<sup>th</sup> in GII (Table 5). Results of MCH, MCV, and MCHC values showed no significant changes in all groups (Table 5).

The result of WBC count showed no significant changes in the level of WBC in all groups (Table 5). Similar result was recorded by (37) who used different dosages of *C. procera* for treatment rabbits coccidiosis, and revealed significant increase ( $P \leq 0.05$ ) at red blood cell count and concentration of hemoglobin attributed to polycythemia, at a high dose of *C. procera*. This result which confirmed by (38), who stated that polycythemia may be due to generalized tissue hypoxia GOT (AST) significantly increased in those of G I and G II during 4<sup>th</sup> and 8<sup>th</sup> weeks, while in GIII only during 8<sup>th</sup> weeks. the highest level was in G I followed by these in G III and G II (Table 6). The serum of GPT (ALT) significantly increased in G II during 4<sup>th</sup> weeks, while in 8<sup>th</sup> were significantly elevated in G I and G II, the highest level was in GI (Table 6). ALP level significantly increased in these of group I,II,III and IV during 4<sup>th</sup> and 8<sup>th</sup> weeks, with the highest levels were in 8<sup>th</sup> in group I then group II (Table 6). Increase in the levels of GOT, GPT and ALP is usually a signal of liver damage (39, 40). These elevations perhaps caused by damage or may be dead of hepatocyte under the venomous impact of active secondary metabolite of plant extract, as a result, this enzyme diffused from the intracellular place (41,42).

In current study, raised of biochemical parameters (GOT, GPT and ALT) of liver assured the toxic effects of the *C. procera* aqueous extract .In contradiction of these results in current study, (10)

observed that, administration of high doses of aqueous extract of *Calotropis procera* not cause significant changes in the activity of AST and ALP and were found significant diminution ( $P \leq 0.05$ ) in the level of ALT in smaller rabbits. Also, Ali, (43) revealed lowering in GOT and GPT after fifteen days at exposure to extract or of this plant. shown in Table (7), results of serum levels of urea and creatinine changes in rabbits of different groups. A non- significant difference ( $P \geq 0.05$ ) was noticed in serum urea levels in all treated groups I, II, III and IV than control group(V) at the 4<sup>th</sup> weeks. Urea significantly elevated ( $P \leq 0.05$ ) in treated groups (I, II, III and IV) during 8<sup>th</sup> weeks in comparison to control group. Creatinine in current study significantly increased in GI and GII during 8th weeks (Table 7). These results go in harmony with the conclusion of (10) who revealed, remarkable raise in urea and creatinine values after treated with aqueous extract of *Calotropis procera*. The Creatinine are removed from plasma via glomerular filtration, then excrete into the urine. Elevated in creatinine values is a signal of renal

dysfunction (44), this injury may be caused by cumulating of effective substance of the aqueous extract of *C. procera* in kidneys (10), tubular epithelial cells may be damaged due to accumulation of toxic substance, or this accumulated substance may cause necrosis in to cells of the renal convoluted tubules (31,45). Impaired the secretary function of the kidney perhaps due to reduced renal blood flow correlating with elevated serum urea level (46). Kidneys damage can be recognized by measure of creatinine level, it is high specific signal to kidney harms (10), other investigators assured the hurtful influence of the *Calotropis procera* extracts on the kidney (13,31, 47). The conclusion, the aqueous extracts of *C. procera* leaves exhibit a toxic effect to rabbits experimentally exposed to it. The effects were a dose dependence in addition to duration of exposure. The dose 162.4 mg/kg b.w. of extract can be considered as sub chronic, as it allows as to fallow the causes especially in heart, liver, and kidney.

**Table 3. Clinical signs that appeared in rabbits used in study**

Signs	Groups				
	I	II	III	IV	V
Depression	+++	++	+	-	-
Diarrhea	+	+	±	-	-
Chewing movement of mouth	+++	++	+	±	-
Engorgement blood vessels at the ear	+++	++	+	-	-
Sneezing	+++	++	±	-	-
Bradycardia	++	+	±	-	-
Convulsion	++	+	-	-	-
Aching of ear and leg	+++	++	+	-	-
Loss of appetite	++	+	±	-	-
Abortion	+	+	-	-	-

+ =signs, += low, += medium, +++=high, ±sometime, - = no signs

**Table 4. Influence of oral administration of different doses of *Calotropis procera* aqueous extract on the body weight, temperature, heart and respiratory rates of rabbits used in the study**

Parameter	Period (weeks)				
	0	2	4	6	8
<b>B.W</b>					
I	1.39±0.07	1.36±0.05	1.36±0.01	1.34±0.05	1.32±0.005
II	1.29±0.03	1.33±0.05	1.35±0.06	1.38±0.08	1.37±0.08
III	1.36±0.004	1.38±0.03	1.42±0.05	1.41±0.04	1.42±0.03
IV	1.22±0.07	1.27±0.10	1.29±0.1	1.29±0.08	1.3±0.06
V	1.26±0.12	1.28±0.11	1.29±0.14	1.31±0.13	1.36±0.11
<b>T</b>					
I	38.8±0.34	38.62±0.26	39.16±0.09	38.44±0.10	38.3±0.36
II	38.42±0.16	38.86±0.22	38.7±0.12	38.48±0.1	38.28±0.28
III	38.78±0.21	38.93±0.15	38.98±0.13	38.7±0.12	38.38±0.11
IV	38.73±0.13	38.3±0.3	39.13±0.30	38.58±0.21	38.85±0.28
V	38.5±0.31	38.98±0.26	38.35±0.18	38.58±0.23	38.58±0.21
<b>H.R</b>					
I	188.6±4.56 <sup>Abc</sup>	175±7.58 <sup>Ab</sup>	171±5.46 <sup>Ab</sup>	151.2±7.29 <sup>Ab</sup>	146±6.44 <sup>Aa</sup>
II	191.2±6.3 <sup>Abc</sup>	183.6±8.13 <sup>Ab</sup>	178.8±6.25 <sup>Bb</sup>	174±6.5 <sup>Bb</sup>	165±4.48 <sup>Ba</sup>
III	189.25±6.1 <sup>Ab</sup>	187.5±6.14 <sup>Bb</sup>	188.25±8.23 <sup>BCb</sup>	184.75±5.04 <sup>BCa</sup>	177.25±5.11 <sup>BCa</sup>
IV	191.5±6.86 <sup>Aa</sup>	190.25±5.11 <sup>Ba</sup>	193.5±4.94 <sup>BCa</sup>	191.75±4.55 <sup>BCDa</sup>	189.75±4.52 <sup>BCDa</sup>
V	191±6.75 <sup>Aa</sup>	190.25±5.5 <sup>Ba</sup>	188.25±6.43 <sup>BCa</sup>	191.75±5.44 <sup>BCDa</sup>	192.25±5.79 <sup>BCDa</sup>
<b>R.R</b>					
I	111.6±6.10 <sup>Ab</sup>	104.4±5.57 <sup>Aa</sup>	101.8±3.63 <sup>Aa</sup>	121.2±6.33 <sup>Bbc</sup>	126.33±4.98 <sup>BCbc</sup>
II	113.4±5.68 <sup>Aa</sup>	111.4±6.32 <sup>Aa</sup>	113±5.43 <sup>Ba</sup>	116.2±7.51 <sup>Aa</sup>	119.8±7.25 <sup>Ba</sup>
III	113.25±6.97 <sup>Aa</sup>	112.5±5.56 <sup>Aa</sup>	107.75±5.65 <sup>Aa</sup>	113.75±6.07 <sup>Aa</sup>	114±6.42 <sup>Aa</sup>
IV	111.75±4.5 <sup>Aa</sup>	108.25±5.12 <sup>Aa</sup>	110.25±5.57 <sup>Ba</sup>	112.25±6.87 <sup>Ba</sup>	110.5±6.5 <sup>Aa</sup>
V	112.25±6.39 <sup>Aa</sup>	108±6.42 <sup>Aa</sup>	112.75±4.97 <sup>Ba</sup>	109±4.81 <sup>Aa</sup>	110.25±4.89 <sup>Aa</sup>

B.W= body weight, T= temperature, H.R= heart rate, R.R= respiratory rates

Results are expressed as mean±SEM. Values with different superscripts along a column are statistically different ( $P \leq 0.05$ ). (5 rabbits in each group)

**Table 5. Influence of oral administration of different doses of *Calotropis procera* aqueous extract on hematological parameters in rabbits used in the study**

Parameter	Period (weeks)		
	0	4W	8W
<b>RBC</b>			
I	6.36±0.13 <sup>Aa</sup>	6.93±0.49 <sup>Aa</sup>	7.17±0.06 <sup>Bb</sup>
II	6.53±0.62 <sup>Aa</sup>	6.69±0.52 <sup>Aa</sup>	6.83±0.03 <sup>Ba</sup>
III	6.52±0.48 <sup>Aa</sup>	6.52±0.17 <sup>Aa</sup>	6.51±0.35 <sup>Aa</sup>
IV	6.27±0.28 <sup>Aa</sup>	6.47±0.18 <sup>Aa</sup>	6.55±0.21 <sup>Aa</sup>
V	6.15±0.21 <sup>Aa</sup>	6.27±0.21 <sup>Aa</sup>	6.26±0.23 <sup>Aa</sup>
<b>Hb</b>			
I	10.76±0.44 <sup>Aa</sup>	11.88±0.48 <sup>Bb</sup>	11.6±0.35 <sup>Ba</sup>
II	10.62±0.42 <sup>Aa</sup>	10.92±0.41 <sup>Aa</sup>	11.7±0.29 <sup>BCb</sup>
III	10.63±0.85 <sup>Aa</sup>	10.9±0.60 <sup>Aa</sup>	10.6±0.53 <sup>Aa</sup>
IV	11.05±0.48 <sup>Aa</sup>	11.45±0.55 <sup>Aa</sup>	11.18±0.41 <sup>Ab</sup>
V	10.98±0.18 <sup>Aa</sup>	11.05±0.26 <sup>Aa</sup>	11.23±0.15 <sup>Ba</sup>
<b>PCV</b>			
I	32±1.00 <sup>Aa</sup>	34±1.05 <sup>Bb</sup>	34±1.16 <sup>Bb</sup>
II	31.2±1.20 <sup>Aa</sup>	32±1.27 <sup>Aa</sup>	34.5±0.87 <sup>Bb</sup>
III	32.25±0.48 <sup>Aa</sup>	32.33±1.08 <sup>Aa</sup>	32.5±0.96 <sup>Aa</sup>
IV	31.75±0.95 <sup>Aa</sup>	32.5±0.65 <sup>Aa</sup>	32.5±0.65 <sup>Aa</sup>
V	31.5±0.65 <sup>Aa</sup>	32±0.82 <sup>Aa</sup>	32±0.41 <sup>Aa</sup>
<b>MCH</b>			
I	16.95±0.73	17.49±1.41	17.03±2.51
II	17.01±2.11	16.86±1.91	17.13±0.5
III	16.40±1.23	16.68±0.51	15.87±1.17
IV	17.12±0.85	17.7±0.84	17.71±0.999
V	17.91±0.69	17.67±0.42	18.18±0.40
<b>MCV</b>			
I	50.39±1.63	50.25±4.50	49.85±7.15
II	50.01±6.27	50.27±5.59	50.52±1.48
III	50.16±3.01	49.60±1.05	48.80±3.76
IV	49.69±0.58	50.23±0.83	50.75±0.8
V	51.33±1.15	51.15±1.06	51.24±1.33
<b>MCHC</b>			
I	33.58±0.40	34.92±0.71	34.12±0.14
II	34.03±0.13	34.13±0.15	33.91±0.01
III	32.86±2.25	33.68±1.32	32.58±0.98
IV	34.89±1.77	35.26±1.75	34.43±1.47
V	34.86±0.59	34.55±0.65	35.08±0.14
<b>WBC</b>			
I	4.69±0.14	4.81±0.15	4.69±0.57
II	4.39±0.14	4.75±0.32	4.62±0.11
III	4.26±0.28	4.47±0.14	4.27±0.32
IV	4.8±0.17	4.76±0.38	4.38±0.1
V	4.28±.28	4.24±0.21	4.3±0.02

Results are expressed as mean±ESM Values with different superscripts along a column are statistically different ( $P \leq 0.05$ ) (5 rabbits in each group)

**Table 6. Influence of oral administration of different doses of aqueous extract of *Calotropis procera* leaves on serum GOT (IU/L), GPT (IU/L), and ALP level (IU/L) in rabbits used in the study**

Parameter	Period (weeks)		
	0	4 week	8 week
<b>GOT</b>			
I	40.52±1.11 <sup>Aa</sup>	62.52±1.1 <sup>Bb</sup>	72.08±0.96 <sup>BCbc</sup>
II	42.48±1.77 <sup>Aa</sup>	62.048±1.57 <sup>Bb</sup>	65.98±1.58 <sup>Bbc</sup>
III	41.87±5.02 <sup>Aa</sup>	53.39±9.2 <sup>Aa</sup>	69.84±7.71 <sup>Bb</sup>
IV	40.15±5.43 <sup>Aa</sup>	49.71±6.81 <sup>Aa</sup>	46.76±4.13 <sup>Aa</sup>
V	42.11±4.13 <sup>Aa</sup>	49.12±5.26 <sup>Aa</sup>	44.39±4.22 <sup>Aa</sup>
<b>GPT</b>			
I	32.39±0.43 <sup>Aa</sup>	31.57±0.97 <sup>Aa</sup>	54.17±1.11 <sup>BCb</sup>
II	31.3±2.31 <sup>Aa</sup>	35.16±2.89 <sup>Bb</sup>	41.21±3.07 <sup>Bbc</sup>
III	33.72±1.83 <sup>Aa</sup>	36.59±2.15 <sup>Ba</sup>	32.64±2.96 <sup>Aa</sup>
IV	32.89±3.48 <sup>Aa</sup>	36.72±2.88 <sup>Ba</sup>	33.05±3.8 <sup>Aa</sup>
V	33.77±2.42 <sup>Aa</sup>	34.99±2.47 <sup>Ba</sup>	34.87±3.99 <sup>Aa</sup>
<b>ALP</b>			
I	159.99±7.73 <sup>Aa</sup>	297.85±4.88 <sup>BCb</sup>	313.25±10.08 <sup>BCDbc</sup>
II	156.54±11.88 <sup>Aa</sup>	310.7±6.34 <sup>BCDb</sup>	303.55±11.10 <sup>BCDb</sup>
III	164.26±8.46 <sup>Aa</sup>	190.7±4.45 <sup>Bb</sup>	243.51±8.4 <sup>BCbc</sup>
IV	160.87±6.91 <sup>Aa</sup>	193.75±4.45 <sup>Bb</sup>	187.69±11.41 <sup>Bb</sup>
V	164.49±7.92 <sup>Aa</sup>	172.23±8.16 <sup>Aa</sup>	169.7±5.47 <sup>Aa</sup>

Results are expressed as mean±SEM. Values with different superscripts along a column are statistically different ( $P \leq 0.05$ ) (5 rabbits in each group)

**Table 7. Influence of oral administration of different doses of aqueous extract of *Calotropis procera* leaves on serum urea level (mg/dl) and serum creatinine (mg/dl) in rabbits used in the study**

Parameter	Period (weeks)		
	0	4week	8week
<b>Urea</b>			
I	28.88±1.99 <sup>Aa</sup>	26.68±2.47 <sup>Aa</sup>	49.29±2.45 <sup>BCDb</sup>
II	27.46±0.89 <sup>Aa</sup>	28.64±1.47 <sup>Aa</sup>	43.35±1.41 <sup>BCb</sup>
III	28.12±1.5 <sup>Aa</sup>	29.47±1.16 <sup>Ba</sup>	34.02±0.98 <sup>Bb</sup>
IV	27.99±0.72 <sup>Aa</sup>	27.12±1.59 <sup>Aa</sup>	31.77±1.33 <sup>Bb</sup>
V	27.48±1.35 <sup>Aa</sup>	27.29±0.91 <sup>Aa</sup>	27.88±1.23 <sup>Aa</sup>
<b>Creatinine</b>			
I	0.48±0.047 <sup>Aa</sup>	0.52±0.037 <sup>Aa</sup>	0.79±0.036 <sup>BCb</sup>
II	0.52±0.057 <sup>Aa</sup>	0.54±0.068 <sup>Aa</sup>	0.87±0.256 <sup>BCDb</sup>
III	0.50±0.079 <sup>Aa</sup>	0.51±0.043 <sup>Aa</sup>	0.60±0.053 <sup>Ba</sup>
IV	0.48±0.014 <sup>Aa</sup>	0.47±0.065 <sup>Aa</sup>	0.50±0.053 <sup>Aa</sup>
V	0.51±0.069 <sup>Aa</sup>	0.49±0.013 <sup>Aa</sup>	0.52±0.039 <sup>Aa</sup>

Results are expressed as mean±SEM Values with different superscripts along a column are statistically different ( $P \leq 0.05$ ). (5 rabbits in each group)

### Acknowledgments

The authors are thankful to the dean of College of Veterinary Medicine, University of Baghdad for providing necessary facilities.

### Conflict of Interest

The authors declare that there is no conflict of interest.

### References

1. Abatan M, Arowolo RO. Toxicity of *Eugenia.uniflora* to Rats. Nig. J. Animal Production.1989;16: 16-19.
2. Parsons WT, Cuthbertson EG. *Noxious Weeds of Australia*. 2nd ed. CSIRO Publishing: Melbourne ;2001.712p.
3. Orwa C, Mutua A, Kindt R, Jamnadass R, Antony S. Agroforestry database: a tree reference and selection guide version 4.0. World Agroforestry Center, Kenya. 2009:1.
4. Quazi S, Kumkum M, Arora S. *Calotropisprocera*: an overview of its phytochemistry and pharmacology. Indian J of Drugs. 2013; 1(2): 63-9.
5. Parrotta JA. Healing plants of Peninsular India. CAB International, Wallingford, UK and New York.2001;944p.
6. Shahat MA, Shihata AMA. Evaluation of the toxicological effects manifested after long term administration of aqueous *Calotropis proceraplant* extract in male and female rabbits. Egypt. J. Hospital Med. 2012; 47: 291-300.
7. Al-Snafi A E. The Constituents and Pharmacological Properties of *Calotropis procera* - An Overview. Int J of Pharm Rev Res. 2015;15: 259-275.
8. Imosemi IO. Evaluation of the toxicity, medicinal use and pharmacological actions of *Calotropis procera*. *EJPMR*.2016;3(9): 28-36.
9. Murti Y, Yogi B, Pathak D. Pharmacognostic standardization of leaves of *Calotropis procera* (Ait.) R. Br. (Asclepiadaceous). Int J Ayurveda Res.2010; 1 (1): 14-17.
10. Pouokam GB, Ahmed H, Dawurung C, Atiku A, David S, Philipe O. Influence of age on sub-chronic toxicity of the aqueous extract of the leaves of *Calotropis procera* on rabbits. J Toxicolo and Envi. Health Sci.2011; 3(5): 119-126.
11. De Lima JM, De Freitas FJ, Amorim RN, Camara AC, Batista JS, Soto-Blanco B. Clinical and pathological effects of *Calotropis procera* exposure in sheep and rats. *Toxicon*. 2011;57 (1): 183-185.
12. Parihar GA, Sharma S, Ghule P, Sharma P, Deshmukh N, Srivastava DN. Anti-inflammatory effect of *Calotropis procera* root bark extract. *Asian J Pharm Life Sci*.2011;1 (1): 29-44.
13. Ahmed OM, Fahim HI, Boules MW, Ahmed HY. Cardiac and testicular toxicity effects of the latex and ethanolic leaf extract of *Calotropis procera* on male albino rats in comparison to abamectin. *Springer Plus*. 2016;5(1): 1644-1665.
14. Basak SK, Bhaumik A, Mohanta A, Singhal P. Ocular toxicity by latex of *Calotropis procera* (Sodom apple). *Indian J. Ophthalmol*. 2009;57(3): 232-234.
15. Lin Z Will Y. Evaluation of drugs with specific organ toxicities in organ-specific cell lines. *Toxico Sci*. 2012;126(1): 114-127.
16. Mohammed A, Ibrahim S, Bilbis LS. Toxicological investigation of aqueous leaf extract of *Calotropis procera* (Ait.) R. Br. in Wister albino rats. *Afr J of Bioch Res*. 2012; 6(7): 90-97.
17. Dixon WJ. Efficient analysis of experimental observation. *Ann. Rev. Pharmacol. Toxicol*. 1980; 20: 441-462.
18. Coles EH. *Veterinary Clinical Pathology* 4th Ed. W.B. Saunders Company, Philadelphia, London and Toronto. 1986; 305-326.
19. Odebiyi A, Sofowora AE. Phytochemical screening of Nigerian medicinal plants. *Lloydia*.1978; 41(3): 234-246.
20. Harborne J B. *Phytochemical methods a guide to modern techniques of plant analysis*.2nd ed. Chapman and Hill, London. 1984.
21. Al-Bid MR. Zurrzus ammamsetur unger abschla B membrane in phoenix dactylifera. Wurzburg University. Wurzburg F.R. of Germany. 1985.
22. Trease GE, Evans MD. *A textbook of Pharmacognosy*, 13th ed. Baillier, Tindal and Caussel, London: 1989;176-180 p.
23. Jaffer HJ, Mahmood MJ, Jawad AM. Phytochemical and biological screening of some Iraqi plant, *Fitoterapia*, LIX. 1983:229.
24. Gomez KA, Gomez AA. *Statistical Procedures for Agricultural Research*. 2nd Ed. John Wiley & Sons, Inc. New York. 1984.
25. Verma R, Satsangi GP, Shrivastava JN. Analysis of phytochemical constituents of the ethanolic and chloroform extracts of *Calotropis procera* using gas chromatography - mass spectroscopy

- (GC-MS) technique. J Med Plants Res. 2013;7(40): 2986-2991.
26. Jato D M, Jacob K A, Zainab A, Okewole P, Atiku A, Shamaki D, *et al.* Effects of the aqueous extract of fresh leaves of *Calotropis procera* on haematological and biochemical parameters in female rabbits. Afr. J. Biotechnol. 2009; 8 (19): 5071-5075.
  27. Al-Robai AA, Abo-Khatwa AN, Danish EY. Toxicological studies on the latex of the usher plants *Calotropis procera* (Alt). R. Br. In Saudi Arabia. III Effects of usher latex on the fine structures, oxygen consumption and  $\text{Na}^{++}\text{K}^{++}$  Transporting AtPase Activity of Albino Rat kidneys. Arab Gulf J of Sci Res.1993; 11(3): 441-455.
  28. Lingrel JB. The physiological significance of the cardiotoxic steroid/ouabain-binding site of the  $\text{Na,K-ATPase}$ . Annu Rev Physiol.2010; 72: 395–412.
  29. Neshet M, Shpolansky U, Rosen H, Lichtstein D. The digitalis like steroid hormones: new mechanisms of action and biological significance. Life Sci.2007; 80: 2093–2107.
  30. Alkhayyat AA, Salih R A. Toxic effects of subacute exposure to *Nerium oleander* leaves hexane extract on the heart of rabbits. Iraqi J Vet. Med;.2017 41(2):54-60.
  31. Mahmoud OM, Adam SEI, Tartour G. The effects of *Calotropis procera* on small ruminants. I. Effects of feeding sheep with the plant. J. Comp. Pathol. 1979; 89: 241–250.
  32. Botha CJ, Penrith ML. Poisonous plants of veterinary and human importance in South Africa. J Ethnopharmacol. 2008; 119: 549-558.
  33. Osbourn A. Saponins and plant defence – a soap story. Trends plant Sci. 1996;1: 4-9.
  34. Faye B. A contribution to the study of the toxicity of *Calotropis procera*. Effect of diets based on *Calotropis procera* on embryonic and neonatal mortality in laboratory mice. Rev Elev Med Vet Pays Trop. 1985; 38 (1): 72–75.
  35. Prakash AO, Gupta RB, Mathur R. Effect of oral administration of two indigenous plant extracts on early and late pregnancy in albino rats. Probe.1978; 17:315-23.
  36. Hilal SH, Youngken JR HW. Certain poisonous plants of Egypt. A Scientific Manual Pharmaceutical Society of Egypt, National Information and Documentation Centers Dokki, Cairo, Egypt. Book.1983: 11.
  37. Seddek A Sh, El-Ghoneimy A A, Dina MW, El-hamd S, Mahmoud E G. Effect of *Calotropis procera* on Rabbits Coccidiosis. Egypt. J. Chem. Environ. Health.2015; 1 (1):768-784.
  38. Evans GO. Animal hemato-toxicology a practical guide for toxicologists and biomedical researchers. CRC Press. New York. 2008.
  39. Chavda R, Validia KR, Gokani R. Hepatoprotective and antioxidant activity of root bark of *Calotropis procera*.br (asclepediaceae). Int. J. Pharmacol. 2010; 6(6): 937-43.
  40. Zhang Y, Huang Y, Deng X, Xu Y, Gao Z, Li H. Iron overload-induced rat liver injury: Involvement of protein tyrosine nitration and the effect of baicalin. Eur J Pharmacol. 2012; (1-3):95-101.
  41. Tilikian SM, Conover M, Tilikian AG. Clinical implications of laboratory tests. The C.V. Mosby Company, Louis, Toronto, London; 1983.
  42. Jimoh FO, Odutuga AA. Changes in the activities of some diagnostic enzymes in some rat tissues following the consumption of thermally oxidized groundnut oil. Nig J Biochem Mol Biol 2001;16: 173-6.
  43. Ali AM. Teratogenic effect and biochemical changes to some rodenticides on certain rodent species. [dissertation]. Cairo, Egypt. University of Al-Azhar: 2006.
  44. Smith CL, Hampton EM. Using estimated creatinine clearance for individualizing drug therapy: a reassessment. DICP.1990; 24:1185-1190.
  45. Parke DV. The Handbook of Environmental Chemistry: 1 Springer Verlag, Berlin; 1982: 141-178.
  46. Whealton A, Watson AJ, Rock RC. Colorimetric determination of serum urea concentration. In Tietz Textbook of Clin Chem Burtis CA, Ashwood ER (eds.). WB Saunders Company London; 1994: 1528-1531.
  47. Jato D M, Jacob K, Zainab A, Okewole P, Atiku A, Shamaki DI, *et al.* Histopathological changes in female rabbits administered with aqueous extract of *Calotropis procera*. Afr. J. Biotechnol. 2016; 15(12): 458-464.

## التأثيرات السمية للمستخلص المائي لأوراق كالتروبيس بروسيرا في الأرناب المسمومة تجريبياً

احمد حنش الزهيري<sup>1</sup>, جنان محمود خلف<sup>2</sup>, صلاح نوري ابراهيم<sup>3</sup>

<sup>1</sup> فرع الطب الباطني, كلية الطب البيطري, جامعة ديالى, ديالى<sup>2</sup>, فرع الطب الباطني, كلية الطب البيطري, جامعة بغداد, بغداد<sup>3</sup>, فرع الفلسفة والادوية, كلية الطب البيطري, جامعة ديالى, ديالى, العراق

### الخلاصة

كالتروبيس بروسيرا سجلت كنبات طبي سام للإنسان والحيوان. التأثيرات السمية للمستخلص المائي للأوراق قيمت تجريبياً في الأرناب. تم حساب الجرعة النصف قاتلة LD<sub>50</sub> (2435.25 ملغم / كغم) باستخدام طريقة الصعود والهبوط. خمسة وعشرون من الأرناب المحلية، بعمر 1-2 سنة، وزن 1-1.9 كغم. قسمت عشوائياً إلى خمس مجموعات، خمسة أرناب لكل منها. عرضت المجموعات الأولى والثانية والثالثة والرابعة إلى 10/1 من الجرعة السامة الوسطية (234.5 ملغم / كغم)، 1 / 12.5 من الجرعة السامة الوسطية (194.8)، 1/15 من الجرعة النصف قاتلة (162.4)، 1/20 من الجرعة نصف القاتلة (121.8 ملغم / كغم من وزن الجسم) من المستخلص وعلى التوالي لمدة 8 أسابيع، في حين المجموعة الخامسة تركت دون التعرض كمجموعة سيطرة. القلوب، الصابونين، العفص، كليكوسيدات القلب، الستيرويد، الكليكوسيدات، التربينويدات والفلافونويد حددت في فحص المواد الكيميائية النباتية. كانت المعايير المعتمدة هي العلامات السريرية التي أظهرتها الحيوانات أثناء الدراسة بالإضافة إلى بعض المعايير الدموية (عدد كرات الدم الحمراء، تركيز الهيموغلوبين، حجم الخلية المرصوص). التغييرات الرئيسية التي لوحظت أثناء مراقبة الحيوانات كانت مضغ الفم، بطء القلب، احتقان الأوعية الدموية، السعال والاكنتاب خاصة في المجموعة 1، 2، 3، انخفاض وزن الجسم في المجموعة الأولى، الاجهاض في المجموعة الأولى والثانية. أظهرت المعايير الدموية والكيميائية الحيوية المعتمدة زيادة كبيرة أثناء الدراسة، ومن هنا يمكن أن نستنتج أن المستخلص المائي لأوراق النبات له تأثير ضار في الأرناب التي اجريت عليها التجربة.

الكلمات المفتاحية: كالتروبيس بروسيرا، الأرناب، السمية، معايير الدم والكيمياء الحيوية